

Older than 12 months Younger than 12 months 1 dose (For unvaccinated TST- or IG-RA-negative older children, adolescents 1 dose and adults from settings with high inci-(if not given earlier) **BCG** dence of TB and/or high leprosy burden and those moving from low to high TB incidence/leprosy burden settings) *If child presents at 3 doses as Pentavalent if age is less <2 months give Hep B single than 6 years. dose f/b Penta 3 doses with minimum interval of 4 weeks If >6yrs give Hepatitis B separately as **Hepatitis B** 3 doses with 4 weeks between 1st and *If child present at ≥2months 2nd dose and minimum 6 months of age start with Pentavalent between 1st and 3rd dose (0,1,6 months) 3 dose series with minimum interval of 4 weeks Polio OPV 3 doses 3 doses **Polio IPV** 1 doses 1 doses **3 doses** with intervals of 2 months between 1st, 2nd dose and 3rd dose after 6 months **Pentavalent** from 1st dose (0,1,6 months) (DPT + Hep B + Hib) 3 doses (If 6-7years of age use TD, if >7 years use Td) 1 dose (if less than 18 months) Measles /MR 1 doses Use MMR 2 doses 4 weeks apart (if age more than 18 months and not received measles) 1 dose of MMR, (if received measles) **MMR** N/A 2 doses of MMR at 4 weeks interval (if not received measles by age 18months) 1 dose (If 6-7 years of age use TD, **DPT Booster** N/A if >7 years use Td) 2 doses of HPV at 6 months interval (if

If interrupted primary series, resume without repeating previous dose

HPV

N/A

If you have any queries, please contact National Immunization Program / Health Protection Agency Phone: 7212232 or 3014333 / Fax: 3014484/ Email: immunization495@health.gov.mv

November 2022

below 15 years) 3 doses at 0,1,6months

(if above 15 years)