

| | Diseases Preventable | Number of doses | Schedule age of 1st dose |
|--|---|---------------------------|---|
| Rota Virus | Diarrhea | 2 months | <ul style="list-style-type: none"> The first dose before 15 weeks of age and the last dose before 6 months of age Rotarix: 2 doses 2 months apart Rotateq: 3 doses 2 months apart |
| Pneumococcal | Pneumonia, meningitis, ear and sinus infection | 2 months | <ul style="list-style-type: none"> 3 doses 2 months apart with or without booster at 15-18 months of age If Initiating vaccination after 12 months of age; Synflorix: between 1 to 5 years of age, 2 doses 2 months apart Prevnar-13: if older than 1 year but less than 2 years old, 2 doses 2 months apart. If older than 2 years, 1 dose |
| Haemophilus Influenzae Type B (Hib Booster) | Pneumonia, meningitis, sinus and ear infection | 12-15 months | <ul style="list-style-type: none"> 1 booster dose Especially for those at high risk of complications such as those with: <ul style="list-style-type: none"> - anatomical /functional asplenia - thalassemia major - immunodeficiency, etc |
| Influenza | Flu, pneumonia, heart problems, ear and sinus infection etc | 6 months of age and above | <ul style="list-style-type: none"> If younger than 9 years of age, 2 doses 4 weeks apart in the first year of vaccination, followed by 1 dose every year If older than 9 years of age or an adult, 1 dose every year Annual vaccination is recommended for health care workers and especially for those at high risk of influenza related complications including people who <ul style="list-style-type: none"> - are pregnant - are thalassemia majors - have chronic heart, lung, liver, kidney disease - are younger than 5 years or older than 65 years - are immunocompromised - have diabetes mellitus - have asthma |
| Chicken pox | Chicken pox | Older than 1 year | <ul style="list-style-type: none"> 2 doses Routine: 1st dose at 12-15months followed by 2nd dose at 4-6 years For children between 16 months and 13 years, 2 doses at least 3 months apart. If older than 13 years of age and not vaccinated/never had chicken pox disease; 2 doses 1 month apart. - Especially recommended for health care workers and childhood educators/care takers |
| Hepatitis A | Hepatitis | Older than 1 year | <ul style="list-style-type: none"> 2 doses, 6 months apart (Note: if older than 19 years, requires adult formulation) |